



TITLE:

# Isolation of (—)S-Allyl-L-cysteine from Garlic

AUTHOR(S):

Suzuki, Tomoji; Sugii, Michiyasu; Kakimoto, Toshio; Tsuboi, Nobuo

---

CITATION:

Suzuki, Tomoji ...[et al]. Isolation of (—)S-Allyl-L-cysteine from Garlic. Bulletin of the Institute for Chemical Research, Kyoto University 1962, 39(6): 416-416

ISSUE DATE:

1962-03-25

URL:

<http://hdl.handle.net/2433/75855>

RIGHT:

## ABSTRACTS

### Isolation of (-)S-Allyl-L-cysteine from Garlic

TOMOJI SUZUKI, MICHIIYASU SUGII, TOSHIO KAKIMOTO and NOBUO TSUBOI

*Chemical and Pharmaceutical Bulletin*, **9**, 251 (1961)

Tracer technique was successfully used for the detection of S-allyl-L-cysteine in garlic. To isolate the new amino acid in large quantities, the neutral garlic amino acids were absorbed on Dowex 50 resin at pH 2.4 and fractionated with 0.2M ammonium formate buffer (pH 3.15). Five kg. of garlic yielded approximately 50mg. of pure S-allyl-L-cysteine, m.p. 218° (decomp.),  $[\alpha]^{20}_D = -8.7^\circ$ . Chemical constants were identical with those of synthesized S-allyl-L-cysteine.